



A complete classification of perfect unitary Cayley graphs of finite rings

Ján Mináč¹, Tung T. Nguyen² and Nguyễn Duy Tân*,³

ABSTRACT

Due to their elegant and simple nature, unitary Cayley graphs have been an active research topic in the literature. These graphs are naturally connected to several branches of mathematics, including number theory, finite algebra, representation theory, and graph theory. In this article, we study the perfectness property of these graphs. More precisely, we provide a complete classification of perfect unitary Cayley graphs associated with finite rings.

ARTICLE HISTORY

Received: 27 October 2025

Revised: 5 January 2026

Accepted: 7 January 2026

Published: 31 March 2026

Communicated by

Eric Andriantiana

<https://doi.org/???>

KEYWORDS

Cayley graphs;

Perfect graphs;

Finite semisimple rings.

MSC

05C25;

05C50;

05C51.

1 Introduction

Let R be a finite unital associative ring. The study of graphs associated with R has received much attention in recent years (see [2, 3, 12] for further discussions on this line of research). One of the first examples of graphs of this type is the family of unitary Cayley graphs, which we will now recall. The unitary Cayley graph on R is defined as the Cayley

*Corresponding author

1. Department of Mathematics, Western University, London, Ontario, Canada N6A 5B7

2. Department of Kinesiology, Western University, London, Ontario, Canada N6A 5B7

3. Faculty of Mathematics and Informatics, Hanoi University of Science and Technology, 1 Dai Co Viet Road, Hanoi, Vietnam

minac@uwo.ca; nguyenthotung@gmail.com; tan.nguyenduy@hust.edu.vn

graph $G_R = \text{Cay}(R, R^\times)$, where R^\times denotes the group of all invertible elements in R . Specifically, G_R is a graph characterized by the following:

1. The vertex set of G_R is R ,
2. Two vertices $a, b \in V(G_R)$ are adjacent if and only if $a - b \in R^\times$.

Various cases of unitary Cayley graphs have been investigated in the literature. To the best of our knowledge, [8] is perhaps the first work that formally introduces the concept of a unitary Cayley graph. In this work, the authors discover some fundamental properties of the unitary graph when R is the ring of integers modulo a given positive integer n , such as their spectra, clique, and independence numbers, planarity, perfectness, and much more. [1] generalizes many results from [8] to the case where R is an arbitrary finite commutative ring. In particular, they are able to classify all perfect unitary Cayley graphs when R is commutative (see [1, Theorem 9.5]). The work [7] extends this line of research to rings that are not necessarily commutative.

Our interest in unitary Cayley graphs stems from the joint work [5], where we classify all prime unitary Cayley graphs amongst other things (see [5, Theorem 4.34]). During our discussions, Sophie Spirkl posed the following question:

Question 1.1. *Can we classify all perfect unitary Cayley graphs?*

We recall that a graph G is said to be perfect if, for every induced subgraph H of G , the chromatic number of H equals the size of its maximum clique. The strong perfect graph theorem gives concrete criteria for a graph to be perfect; namely, a graph is perfect if and only if neither the graph itself nor its graph complement contains an induced odd cycle of length of at least five (see [6]).

The goal of this article is to provide a complete answer to Question 1.1. For the precise statement, we refer readers to Theorem 4.5. We remark that our approach combines experimental and theoretical mathematics. Specifically, we use SageMath to generate unitary Cayley graphs and then utilize algorithms from the Python library NetworkX to identify relevant cycles within these graphs and check whether two given graphs are isomorphic. As it turns out, when they exist these cycles exhibit several common features, enabling us to find a pattern that is applicable to general cases (of course, we need to show that the pattern that we found *actually* works! For that, we need pure mathematics.)

1.1 Code

The code that we developed to generate unitary graphs over a finite ring and do experiments on them can be found at [10].

2 Reduction to the semisimple case

In this section, we use some structure theorem for finite rings to reduce Question 1.1 to the case where R is a semisimple ring. We first start with the following observation. While we

believe that this observation is well-known among the experts, we cannot find a reference for it. For the sake of completeness, we provide our proof here.

Proposition 2.1. *Let R be a finite ring and $r \in R$. Then the following conditions are equivalent.*

1. r is invertible.
2. r is left invertible.
3. r is right invertible.

Proof. By definition, (1) implies (2) and (3). Additionally, (2)+(3) implies (1). Therefore, it is sufficient to show that (2) and (3) are equivalent. We will show that (2) implies (3). The statement that (3) implies (2) can be proved using the same argument.

Now, suppose that (2) holds. Let $r \in R$ such that r is left-invertible; i.e., there exists $s \in R$ such that $sr = 1$. Let us consider the multiplication by r map $m_r: R \rightarrow R$ defined by $m_r(a) = ra$. This is an injective group homomorphism on $(R, +)$. Indeed, let $a \in \ker(m_r)$. Then $0 = s(ra) = (sr)a = a$ and hence $\ker(m_r) = 0$. Because R is a finite set, we conclude that m_r is also surjective. In particular, we can find s' such that $rs' = 1$. This shows that r is right invertible and therefore invertible. \square

Remark 2.2. *By Proposition 2.1, there will be no ambiguity when we write R^\times .*

We recall that the Jacobson radical $\text{Rad}(R)$ of R is the intersection of all left maximal ideals in R (see [11, Chapter 4.3]). It turns out that $\text{Rad}(R)$ is a two-sided ideal in R . By [5, Proposition 4.30], we know that $\text{Rad}(R)$ is a homogeneous set in the unitary Cayley graph G_R . Additionally, by for [5, Corollary 4.2], we know that

$$G_R \cong G_{R^{\text{ss}}} * E_n,$$

here $R^{\text{ss}} = R/\text{Rad}(R)$ is the simplification of R , E_n is the empty graph on $n = |\text{Rad}(R)|$ vertices, and $*$ denotes the wreath product of two graphs (see [5, Definition 2.5] for the definition of the wreath product of graphs).

Remark 2.3. *Technically speaking, [5] only deals with commutative rings. However, the arguments for [5, Theorem 4.30, Corollary 4.2] can be applied directly to all finite rings.*

By the strong graph theorem, we can see that $G_1 * G_2$ is perfect if and only if each G_1 and G_2 is. Since the empty graph E_n is perfect, we have the following immediate consequence.

Proposition 2.4. *G_R is perfect if and only if $G_{R^{\text{ss}}}$ is perfect.*

Therefore, from now on, we can assume that $R = R^{\text{ss}}$; i.e., R is semisimple. By the Artin-Wedderburn structure theorem, we know that

$$R = \prod_{i=1}^s R_i \times \prod_{i=1}^r M_{d_i}(F_i).$$

Here R_i is a finite field such that $2 \leq |R_1| \leq |R_2| \leq \dots \leq |R_s|$. Additionally, $d_i \geq 2$, and F_i is a finite field. We can then see that G_R is a direct product of the unitary Cayley graphs

$$G_R = \prod_{i=1}^s G_{R_i} \times \prod_{i=1}^r G_{M_{d_i}(\mathbb{F}_i)} = \prod_{i=1}^s K_{|R_i|} \times \prod_{i=1}^r G_{M_{d_i}(\mathbb{F}_i)}.$$

The case where R is commutative is treated in [1]. To break down the problem, we will deal with one factor at a time. For these reasons, we will first consider the case where R is a matrix ring over a finite field in the next section.

3 R is a matrix ring.

Let $d \geq 2$ and F be a finite field. Let $M_d(F)$ be the ring of $d \times d$ matrices with coefficients in F . Let $G_{M_d(F)}$ be the unitary Cayley graph on $M_d(F)$. We study the following question.

Question 3.1. *When is $G_{M_d(F)}$ a perfect graph?*

We remark that there is a diagonal embedding $F^d \rightarrow M_d(F)$. Consequently, G_{F^d} is naturally an induced subgraph of $G_{M_d(F)}$. Consequently, if $G_{M_d(F)}$ is perfect, then so is G_{F^d} . By [1, Theorem 9.5], we conclude that either $F = \mathbb{F}_2$ or $d = 2$. In summary, we have just proved the following.

Proposition 3.2. *If $M_d(F)$ is perfect, then either $d = 2$ or $F = \mathbb{F}_2$.*

We remark that when $F = \mathbb{F}_2$, the graph G_{F^d} is bipartite and hence perfect. One can naturally ask whether $G_{M_d(F)}$ is bipartite. The answer is no. To show this, we need to recall the following lemmas.

Lemma 3.3 (See [4, Proposition 2.6]). *Let G be a finite group and S a symmetric subset of G . Suppose further that $\text{Cay}(G, S)$ is connected. Then $\text{Cay}(G, S)$ is bipartite if and only if there exists an index 2 subgroup H of G such that $H \cap S = \emptyset$.*

Lemma 3.4 (See [9]). *If $d \geq 2$ then every matrix in $M_d(F)$ can be written as the sum of two invertible matrices.*

Proposition 3.5. *If $d \geq 2$ then $G_{M_d(F)}$ is not bipartite.*

Proof. Assume that $G_{M_d(F)}$ is bipartite. Then by Lemma 3.3 we can find an additive subgroup H of $M_d(F)$ with index 2 such that $H \cap GL_d(F) = \emptyset$. By this assumption, we conclude that if $a, b \in GL_d(F)$, then $a + b \in H$. By Lemma 3.4, we conclude that $H = M_d(F)$, which is a contradiction. We conclude that $G_{M_d(F)}$ is not bipartite. \square

Using some Sagemath code, we can check that there are no induced odd cycles of length at least 5 in $G_{M_2(\mathbb{F}_2)}$ (see [10] where we do a brute force search for odd-cycles on $G_{M_2(\mathbb{F}_2)}$). However, $G_{M_2(\mathbb{F}_3)}$ contains the following induced 5-cycle $A_1 \rightarrow A_2 \rightarrow A_3 \rightarrow A_4 \rightarrow A_5$, where

$$A_1 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, A_2 = \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}, A_3 = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}, A_4 = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}, A_5 = \begin{bmatrix} 1 & 2 \\ 2 & 2 \end{bmatrix}.$$

We conclude that $G_{M_2(\mathbb{F}_3)}$ is not perfect. More generally, if $\text{char}(F) \neq 2$, we can find the following induced 5-cycle in $G_{M_2(F)}$

$$A_1 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, A_2 = \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}, A_3 = \begin{bmatrix} -2 & -2 \\ -1 & -1 \end{bmatrix}, A_4 = \begin{bmatrix} -1 & -2 \\ -1 & -2 \end{bmatrix}, A_5 = \begin{bmatrix} -2 & -1 \\ -1 & -1 \end{bmatrix}.$$

We conclude the following proposition.

Proposition 3.6. *If $\text{char}(F) \neq 2$ then $G_{M_2(F)}$ contains an induced 5-cycle and hence $G_{M_2(F)}$ is not perfect.*

In the case $\text{char}(F) = 2$ and $F \neq \mathbb{F}_2$, we find the following induced 5-cycle in $G_{M_2(F)}$.

$$A_1 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, A_2 = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, A_3 = \begin{bmatrix} z+1 & 0 \\ 1 & 0 \end{bmatrix}, A_4 = \begin{bmatrix} 1 & z+1 \\ 1 & z+1 \end{bmatrix}, A_5 = \begin{bmatrix} z+1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Here z is any element in $F \setminus \mathbb{F}_2$. We conclude that

Proposition 3.7. *If $\text{char}(F) = 2$ and $F \neq \mathbb{F}_2$ then $G_{M_2(F)}$ contains an induced 5-cycle and hence $G_{M_2(F)}$ is not perfect.*

For the case $M_d(\mathbb{F}_2)$ with $d \geq 3$, we also have

Proposition 3.8. *If $d \geq 3$, then $G_{M_d(F)}$ contains an induced 5-cycle and hence $G_{M_d(\mathbb{F}_2)}$ is not perfect.*

Proof. We will show that the following vertices produce an induced 5-cycle (with some help from computers for small d)

$$0 \rightarrow I + A \rightarrow A \rightarrow I + A + B^T \rightarrow I + B.$$

Here

$$A = \begin{bmatrix} 1 & 1 & \cdots & 0 & 0 & 0 \\ 1 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix}, \text{ and } B = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

In other words, the entries in the upper left 2×2 -minor of A are 1 while the other entries are 0 and B is a Jordan block of size d .

It is straightforward to see that $A^d = B^d = 0$. Hence $I + A$, $I + B$, $(I + A + B^T) - A = I + B^T$ are invertible. We have $|(I + A) - (I + B)| = |A - B| = 0$ since the d th row of $A - B$ is the zero row. We also have $|I + A + B^T| = 0$ since the first column of $I + A + B^T$ is the

zero column. On the other hand, $|(I + B) - A| = 0$ since the first row of $I + B - A$ is the zero row.

Now we only need to show that $(I + A + B^T) - (I + B) = A + B + B^T$ is invertible. For convenience, for each $n \geq 2$ let $D_n = B_n + B_n^T$, where B_n is a Jordan block of size of n . By expanding the determinant of D_n along the first column and then along the first row we get $|D_n| = |D_{n-1}|$. Hence $|D_n| = |D_2| = 1$ if n is even and $|D_n| = |D_3| = 0$ if n is odd.

Now

$$\begin{aligned}
|A + B + B^T| &= \begin{vmatrix} 1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & & & \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 1 & 0 \end{vmatrix} \\
&= \begin{vmatrix} 1 & 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & & \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{vmatrix} \quad (\text{expanding the determinant along the 1st row}) \\
&= |D_{d-2}| - \begin{vmatrix} 1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & & \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{vmatrix} \quad (\text{expanding the determinant along the 1st row}) \\
&= |D_{d-2}| - |D_{d-3}| \quad (\text{expanding the second determinant along the 1st column}) \\
&= 1.
\end{aligned}$$

Hence, $A + B + B^T$ is invertible and we are done. \square

In summary, we have proved the following theorem.

Theorem 3.9. $G_{M_d(F)}$ is perfect if and only if $d = 2$ and $F = \mathbb{F}_2$.

4 The general case

As discussed at the end of 2, for Question 1.1 we can assume that R is semisimple. Furthermore, we can suppose that R has the following structure

$$R = \prod_{i=1}^s R_i \times \prod_{i=1}^r M_{d_i}(F_i).$$

Here, R_i is a finite field such that $0 \leq |R_1| \leq |R_2| \leq \dots \leq |R_s|$. Additionally, $d_i \geq 2$, and F_i is a finite field. With this decomposition, G_R is the following direct product

$$G_R = \prod_{i=1}^s G_{R_i} \times \prod_{i=1}^r G_{M_{d_i}(\mathbb{F}_i)} = \prod_{i=1}^s K_{|R_i|} \times \prod_{i=1}^r G_{M_{d_i}(\mathbb{F}_i)}.$$

If $|R_1| = 2$, then G_R is bipartite, hence perfect. Therefore, we can assume that $|R_i| \geq 3$ for all $1 \leq i \leq s$.

Lemma 4.1. *Suppose R_1, R_2, R_3 are finite fields such that $|R_i| \geq 3$, for all $1 \leq i \leq 3$. Then $G_{R_1} \times G_{R_2} \times G_{R_3}$ contains an induced 5-cycle.*

Proof. Let $a \in R_2 \setminus \{0, 1\}$, $b \in R_3 \setminus \{0, 1\}$ and $c \in R_1 \setminus \{0, 1\}$. Then one can check that

$$(0, 0, 0) \rightarrow (1, 1, 1) \rightarrow (0, a, b) \rightarrow (1, 1, 0) \rightarrow (c, a, 1)$$

is an induced 5-cycle. □

Lemma 4.2. *Let G_1, G_2, \dots, G_d be graphs and $k \geq 2$ a fixed integer. Suppose that for each $1 \leq i \leq d$,*

$$v_{i1} \rightarrow v_{i2} \rightarrow v_{i3} \rightarrow v_{i4} \rightarrow \dots \rightarrow v_{ik}$$

is a closed path of length k in G_i (we do not require that $\{v_{ij}\}_{1 \leq j \leq k}$ are different, except for $i = 1$). Suppose further that the induced graph on $\{v_{11}, v_{12}, v_{13}, \dots, v_{1k}\}$ is a k -cycle. For each $1 \leq j \leq k$, let

$$v_j = (v_{ij})_{1 \leq i \leq d} \in G_1 \times G_2 \times \dots \times G_d.$$

Then the induced graph on $\{v_j\}_{1 \leq j \leq k}$ is an induced k -cycle.

Proof. Clearly v_1, v_2, \dots, v_k are pairwise different and $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k$ is a closed path. Since the induced graph on $\{v_{11}, v_{12}, v_{13}, \dots, v_{1k}\}$ is a k -cycle, we see that if $i \neq j$ then v_{1i} and v_{1j} are not connected, and hence v_i and v_j are not connected as well. □

By our assumption that $3 \leq |R_1| \leq |R_2| \leq \dots \leq |R_s|$, K_3 is always a subgraph of G_{R_i} . By Lemma 3.4, we also know that K_3 is a subgraph of $G_{M_{d_i}(F_i)}$ as well. Since K_3 contains a closed path of length 5, G_{R_i} and $G_{M_{d_i}(F_i)}$ both contain a closed path of length 5. We conclude that if R contains a factor of the form $M_d(F) \neq M_2(\mathbb{F}_2)$ or $s \geq 3$, then G_R is perfect as it will contain an induced 5-cycle. We now deal with the remaining cases, namely

$$R = \prod_{i=1}^s R_i \times (M_2(\mathbb{F}_2))^r,$$

where $s \leq 2$ and $r \geq 0$. For these cases, we have the following observation.

Lemma 4.3. *Suppose that $R = M_2(\mathbb{F}_2) \times F$, where F is a field such that $F \neq \mathbb{F}_2$. Let $\alpha \in F \setminus \mathbb{F}_2$. Then the following elements produce an induced 5-cycle on G_R*

$$r_1 = \left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, 0 \right), r_2 = \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, 1 \right), r_3 = \left(\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}, \alpha \right), r_4 = \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, 0 \right), r_5 = \left(\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \alpha \right).$$

Consequently, G_R is not perfect.

For the case $R = M_2(F_2) \times M_2(\mathbb{F}_2)$ we have the following

Lemma 4.4. *Suppose that $R = M_2(\mathbb{F}_2) \times M_2(\mathbb{F}_2)$. Then the following elements produce an induced 5-cycle on G_R*

$$r_1 = \left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \right), r_2 = \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right), r_3 = \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \right), \\ r_4 = \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \right), r_5 = \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right).$$

Consequently, G_R is not perfect.

In summary, we have the following theorem which classifies all perfect unitary Cayley graphs.

Theorem 4.5. *Suppose that R is a finite ring such that its semisimplification has the following decomposition*

$$R^{ss} = \prod_{i=1}^s R_i \times \prod_{i=1}^r M_{d_i}(F_i).$$

Here, R_i is a finite field such that $2 \leq |R_1| \leq |R_2| \leq \dots \leq |R_s|$. Additionally, $d_i \geq 2$, and F_i is a finite field. Then G_R is a perfect graph if and only if one of the following conditions is satisfied.

1. $|R_1| = 2$. In this case, G_R is a bipartite graph and hence perfect.
2. $s \leq 2$ and $r = 0$. In other words, either $R^{ss} = R_1$ or $R^{ss} = R_1 \times R_2$.
3. $R^{ss} = M_2(\mathbb{F}_2)$.

Acknowledgements

We thank Maria Chudnovsky, Michal Cizek, Logan Crew, Sophie Spirkl, and Torsten Sander for the enlightening discussions about unitary Cayley graphs and related topics. We are especially grateful to Sophie Spirkl for raising Question 1.1. We also thank Sunil Chebolu for introducing us to the article [9]. Last but not least, we would like to thank the referee for their suggestions which improved our exposition.

Funding

JM is partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) grant R0370A01. He gratefully acknowledges the Western University Faculty of Science Distinguished Professorship 2020-2021. TTN is partially supported by an AMS-Simons Travel Grant. NDT is partially supported by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.04-2023.21

References

- [1] R. Akhtar, M. Boggess, T. Jackson-Henderson, I. Jiménez, R. Karpman, A. Kinzel, and D. Pritikin, *On the unitary Cayley graph of a finite ring*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 117, 13 pages.
- [2] D. F. Anderson, T. Asir, A. Badawi, and T. Tamizh Chelvam, *Graphs from rings*, Springer, Cham, 2021.
- [3] T. Asir and T. Tamizh Chelvam, *On the genus of generalized unit and unitary Cayley graphs of a commutative ring*, Acta Math. Hungar. **142** (2014), no. 2, 444–458.
- [4] A. Biswas, *On a Cheeger type inequality in Cayley graphs of finite groups*, European J. Combin. **81** (2019), 298–308.
- [5] M. Chudnovsky, M. Cizek, L. Crew, J. Mináč, T. T. Nguyen, S. Spirkl, and N. D. Tân, *On prime Cayley graphs*, J. Comb. **17** (2026), no. 2, 223-252.
- [6] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, *The strong perfect graph theorem*, Ann. of Math. **164** (2006), 51–229.
- [7] D. Kiani and M. M. Haji Aghaei, *On the unitary Cayley graph of a ring*, Electron. J. Combin. **19** (2012), P10.
- [8] W. Klotz and T. Sander, *Some properties of unitary Cayley graphs*, Electron. J. Combin. **14** (2007), R45.
- [9] N. J. Lord, *Matrices as sums of invertible matrices*, Math. Mag. **60** (1987), 22–24.
- [10] J. Mináč, T. T. Nguyen, and N. D. Tân, *Unitary Cayley graphs over a finite ring*, Available at https://github.com/tungprime/perfect_unitary_cayley, 2024.
- [11] R. S. Pierce, *Associative algebras*, Studies in the History of Modern Science, Vol. 9, Springer-Verlag, New York–Berlin, 1982.
- [12] H. Su and G. Tang, *When unit graphs are isomorphic to unitary Cayley graphs of rings?*, J. Algebra Appl. **24** (2025), no. 11, Paper No. 2550267, 9 pages.